


Equation of State of Neutron-Rich Matter and Implications for Neutron Stars

OHIO
UNIVERSITY

Christian Drischler

14th Conference on the Intersections of Particle and Nuclear Physics 2022

September 1, 2022



BUQEYE Collaboration



Ribbon-cutting ceremony
May 2, 2022

See also Brad Sherrill's talk (Friday):
Status and Prospects with FRIB



Facility for Rare Isotope Beams
at Michigan State University


Samuel L. Stanley
President of MSU

Jennifer M. Granholm
Secretary of Energy

Recent neutron star observations

What is the maximum neutron star mass?

GW170817
GRB170817A
AT2017gfo




+ Virgo (Italy)
+ (KAGRA)

heaviest & fastest known galactic neutron star

$$M = 2.35 \pm 0.17 M_{\odot}$$

J0952-0607: Romani et al. (2022)



NICER
soft X-ray telescope

multi-messenger astronomy

Recent neutron star observations

What is the maximum neutron star mass?

GW170817
GRB170817A
AT2017gfo

heaviest & fastest known galactic neutron star


$$M = 2.35 \pm 0.17 M_{\odot}$$

J0952-0607: Romani et al. (2022)

See also Ben Margalit's talk (this session):
*Mergers, kilonovae, and two-solar-mass neutron stars:
What does astrophysics tell us about the nuclear EOS?*






See also K. Chatzilioannou's talk (Saturday):
GWs from binary neutron stars and neutron star black-hole mergers




multi-messenger astronomy

Structure of cold neutron stars





Ab initio workflow (idealized)



CD & Bogner, Few Body Syst. **62**, 109

Here: nuclear equation of state (EOS)
energy per particle (and derived quantities)


$$\frac{E}{A}(n, \delta, T)$$



theory of strong interactions
QCD is nonperturbative at the low energies
relevant for nuclear physics (cf. pQCD & LQCD)

CD, Haxton, McElvain, Mereghetti *et al.*, PPNP **121**, 103888



Ab initio workflow (idealized)



CD & Bogner, Few Body Syst. **62**, 109

Here: nuclear equation of state (EOS)
energy per particle (and derived quantities)


$$\frac{E}{A}(n, \delta, T)$$



theory of strong interactions
QCD is nonperturbative at the low energies
relevant for nuclear physics (cf. pQCD & LQCD)

CD, Haxton, McElvain, Mereghetti et al., PPNP **121**, 103888

Ab initio workflow (idealized)



Here: nuclear equation of state (EOS)
energy per particle (and derived quantities)

$$\frac{E}{A}(n, \delta, T)$$

baryon density n
neutron excess δ
temperature $T (= 0)$

computational framework

solves the (many-body) Schrödinger equation
requires a nuclear potential as input


chiral effective field theory

provides microscopic interactions consistent with
the symmetries of *low-energy* QCD

theory of strong interactions

QCD is nonperturbative at the low energies
relevant for nuclear physics (cf. pQCD & LQCD)

Ab initio workflow (idealized)



Here: nuclear equation of state (EOS)
energy per particle (and derived quantities)

See also Heiko Hergert's slides:
*A Status Update on
Ab Initio Calculations in Nuclear Physics*

computational framework
solves the (many-body) Schrödinger equation
requires a nuclear potential as input

chiral effective field theory
provides microscopic interactions consistent with
the symmetries of *low-energy* QCD


theory of strong interactions
QCD is nonperturbative at the low energies
relevant for nuclear physics (cf. pQCD & LQCD)

Rigorous UQ for nuclear matter



CD, Furnstahl, Melendez,
Phillips, PRL 125, 202702

OHIO
UNIVERSITY



Chiral Effective Field Theory (nucleons & pions)

dominant approach for deriving *microscopic* interactions
consistent with the symmetries of *low-energy QCD*

three- and four-neutron forces predicted through N³LO

See also Evgeny Epelbaum's slides (LENPIC):
Chiral EFT for low-energy nuclear physics


See also Maria Piarulli's slides:
Analyzing the nuclear interaction: challenges
and new opportunities

Rigorous UQ for nuclear matter



CD, Furnstahl, Melendez,
Phillips, PRL 125, 202702

OHIO
UNIVERSITY




Chiral Effective Field Theory (nucleons & pions)

dominant approach for deriving *microscopic* interactions
consistent with the symmetries of *low-energy QCD*


three- and four-neutron forces predicted through N³LO

enables **uncertainty quantification** (EFT truncation)

Bayesian methods are powerful tools for quantifying &
propagating EFT uncertainties based on *falsifiable* model
assumptions



Open-source software & tutorials (Jupyter): <https://bugeye.github.io>



BUQEYE Collaboration


Bayesian
Uncertainty
Quantification:
Errors for
Your
EFT

Rigorous UQ for nuclear matter



CD, Furnstahl, Melendez,
Phillips, PRL 125, 202702

OHIO
UNIVERSITY



Chiral Effective Field Theory (nucleons & pions)

dominant approach for deriving *microscopic* interactions
consistent with the symmetries of *low-energy QCD*

three- and four-neutron forces predicted through N³LO

enables **uncertainty quantification** (EFT truncation)


Bayesian methods are powerful tools for quantifying &
propagating EFT uncertainties based on *falsifiable* model
assumptions

An example: symmetric matter

$$y = \frac{E}{A}, \quad k = 4 \quad (\text{N}^3\text{LO})$$

Uncertainty bands depict
68% credibility regions

$$y = y_k + \delta y_k$$




Rigorous UQ for nuclear matter



CD, Furnstahl, Melendez,
Phillips, PRL 125, 202702

OHIO
UNIVERSITY



Chiral Effective Field Theory (nucleons & pions)

dominant approach for deriving *microscopic* interactions
consistent with the symmetries of *low-energy QCD*

three- and four-neutron forces predicted through N³LO

enables **uncertainty quantification** (EFT truncation)


Bayesian methods are powerful tools for quantifying &
propagating EFT uncertainties based on *falsifiable* model
assumptions

An example: symmetric matter

$$y = \frac{E}{A}, \quad k = 4 \quad (\text{N}^3\text{LO})$$

Uncertainty bands depict
68% credibility regions

$$y = y_k + \delta y_k$$




Rigorous UQ for nuclear matter



CD, Furnstahl, Melendez,
Phillips, PRL 125, 202702

OHIO
UNIVERSITY



Chiral Effective Field Theory (nucleons & pions)

dominant approach for deriving *microscopic* interactions
consistent with the symmetries of *low-energy QCD*

three- and four-neutron forces predicted through N³LO

enables **uncertainty quantification** (EFT truncation)


Bayesian methods are powerful tools for quantifying &
propagating EFT uncertainties based on *falsifiable* model
assumptions

An example: symmetric matter

$$y = \frac{E}{A}, \quad k = 4 \quad (\text{N}^3\text{LO})$$

Uncertainty bands depict
68% credibility regions

$$y = y_k + \delta y_k$$




Rigorous UQ for nuclear matter



CD, Furnstahl, Melendez,
Phillips, PRL 125, 202702

OHIO
UNIVERSITY



Chiral Effective Field Theory (nucleons & pions)

dominant approach for deriving *microscopic* interactions
consistent with the symmetries of *low-energy QCD*

three- and four-neutron forces predicted through N^3LO

enables **uncertainty quantification** (EFT truncation)


Bayesian methods are powerful tools for quantifying &
propagating EFT uncertainties based on *falsifiable* model
assumptions

An example: symmetric matter

$$y = \frac{E}{A}, \quad k = 4 \quad (N^3LO)$$

Uncertainty bands depict
68% credibility regions

$$y = y_k + \delta y_k$$




Rigorous UQ for nuclear matter



CD, Furnstahl, Melendez,
Phillips, PRL 125, 202702

OHIO
UNIVERSITY



Chiral Effective Field Theory (nucleons & pions)

dominant approach for deriving *microscopic* interactions
consistent with the symmetries of *low-energy QCD*

three- and four-neutron forces predicted through N³LO

enables **uncertainty quantification** (EFT truncation)


Bayesian methods are powerful tools for quantifying &
propagating EFT uncertainties based on *falsifiable* model
assumptions

An example: symmetric matter


$$y = \frac{E}{A}, \quad k = 4 \quad (\text{N}^3\text{LO})$$

Uncertainty bands depict
68% credibility regions

$$y = y_k + \delta y_k$$



Ab initio workflow (idealized)



Here: nuclear equation of state (EOS)
energy per particle (and derived quantities)

$$\frac{E}{A}(n, \delta, T)$$

baryon density n
neutron excess δ
temperature $T (= 0)$

Here: many-body perturbation theory (MBPT)


computationally efficient method (HPC-friendly)
allows to estimate many-body uncertainties

Widely applicable:

- ✓ arbitrary proton fractions
- ✓ finite temperature
- ✓ optical potentials, linear response, nuclei, ...

Other frameworks include quantum Monte Carlo,
coupled cluster, and self-consistent Green's functions

Ab initio workflow (idealized)



Here: nuclear equation of state (EOS)

energy

See also Stefano Gandolfi's slides:
Recent QMC calculations of properties
of nuclei & nuclear matter

A

temperature $T (= 0)$

Here: many-body perturbation theory (MBPT)

computationally efficient method (HPC-friendly)
allows to estimate many-body uncertainties

Widely applicable:

- ✓ arbitrary proton fractions
- ✓ finite temperature
- ✓ optical potentials, linear response, nuclei, ...

Other frameworks include quantum Monte Carlo,
coupled cluster, and self-consistent Green's functions

Renaissance of MBPT

CD, Hebeler, Schwenk, PRL 122, 042501


CD, McElvain *et al.*, in prep.

OHIO
UNIVERSITY



automated approach to MBPT for nuclear matter with NN, 3N, and 4N forces

- implementation of arbitrary diagrams has become straightforward (numerically exact)
- multi-dimensional momentum integrals: improved VEGAS
- GPU-accelerated normal ordering of **3N interactions**
- propagation of importance sampling distributions
- controlled evaluation of 1000s of MBPT diagrams



high-order MBPT
calculations of the EOS

automated code
generation

analytic expressions
interaction & MBPT diagrams

automated diagram
generation

Application to dilute Fermi gas:

Wellenhofer, CD, Schwenk, PRC 104, 014003 & PLB 802, 135247

Arthuis *et al.*,
Comput. Phys. 240, 202

The number of diagrams increases rapidly!

1, 3, 39, 840, 27 300, 1 232 280, ...

$n =$ 2 3 4 5 6 7

Integer sequence A064732:
Number of labeled Hugenholtz diagrams with n nodes.

with automated diagram generation



automated approach
to MBPT for nuclear matter

MBPT: an HPC application



#2 (U.S.)


Summit @ Oak Ridge Leadership Computing Facility

202 752 CPU Cores

27 648 Nvidia GPUs

122.3 peta flops

MBPT: an HPC application




Drischler, McElvain *et al.*, in prep.

#2 (U.S.)

Summit @ Oak Ridge Leadership Computing Facility

202 752 CPU Cores
27 648 Nvidia GPUs
122.3 peta flops

Ab initio workflow (idealized)



nuclear EOS with quantified uncertainties
energy per particle (and derived quantities)

$$\frac{E}{A}(n, \delta, T)$$

baryon density n
neutron excess δ
temperature $T (= 0)$

Uncertainty quantification


robust estimates of theoretical uncertainties using
Bayesian machine learning via Gaussian Processes
uncertainties in EFT-based calculations due to:

- truncating the EFT expansion
- applying many-body (and other) approximations
- fitting LECs to experimental data


First chiral potentials with uncertainties
fully quantified and their applications:

Wesolowski, Svensson *et al.*, PRC **104**, 064001
Djärv, Ekström *et al.*, PRC **105**, 014005


Neutron matter | saturation in symmetric matter



$$\left. \frac{d}{dn} \frac{E}{A}(n, 0) \right|_{n=n_0} = 0$$




Neutron matter | saturation in symmetric matter




CD, Holt, and Wellenhofer, Annu. Rev. Nucl. Part. Sci. **71**, 403

saturation point: **fine-tuned cancellation**
between the kinetic and interaction
contributions (ideal testbed for chiral EFT)



Coester band overlaps with the empirical
box (but limited meaning without errors)
Annotations: (λ / Λ_{3N}) in fm^{-1} or (Λ) in MeV




Piekarewicz & Fattoyev, Phys. Today **72**, 7

needed: improved predictions with
novel NN+3N interactions and
robust uncertainty quantification

Nuclear symmetry energy


$$E_{\text{sym}}(n) \approx \frac{E}{N}(n) - \frac{E}{A}(n)$$




See also Betty Tsang's talk:
Symmetry Energy (Saturday)

Nuclear symmetry energy

$$E_{\text{sym}}(n) \approx \frac{E}{N}(n) - \frac{E}{A}(n)$$



excellent agreement with experiment



$$S_2(n) \equiv S_v + \frac{L}{3} \left(\frac{n - n_0}{n_0} \right) + \dots$$

$$\text{pr}(S_v, L | \mathcal{D}) = \int \text{pr}(S_v, L | \mathcal{D}, n_0) \text{pr}(n_0 | \mathcal{D}) dn_0$$


$$\text{pr}(n_0 | \mathcal{D}) \approx 0.17 \pm 0.01 \text{ fm}^{-3}$$

Correlations are important:
uncertainties can be smaller
than one *might* naively think


CD, Holt *et al.*, ARNPS **71**, 403
Lattimer & Lim, APJ **771**, 51

Nuclear symmetry energy

$$E_{\text{sym}}(n) \approx \frac{E}{N}(n) - \frac{E}{A}(n)$$



excellent agreement with experiment




**See also Juliette Mammei's slides (next talk):
The PREX and CREX Experiments**

$$\text{pr}(S_v, L | \mathcal{D}) = \int \text{pr}(S_v, L | \mathcal{D}, n_0) \text{pr}(n_0 | \mathcal{D}) dn_0$$

$$\text{pr}(n_0 | \mathcal{D}) \approx 0.17 \pm 0.01 \text{ fm}^{-3}$$

Correlations are important:
uncertainties can be smaller than one *might* naively think


$$S_2(n) \equiv S_v + \frac{L}{3} \left(\frac{n - n_0}{n_0} \right) + \dots$$



Reinhard *et al.*, PRL 127, 232501
Reed, Fattoyev *et al.*, PRL 126, 172503
Piekarewicz, PRC 104, 024329

“Tension” between PREX-II and different theoretical approaches at the ~68-95% level

Exploring the limits of chiral EFT




CD, Melendez *et al.*, PRC 102, 054315

Bayesian inference of the
in-medium breakdown scale

**But: at what density does
chiral EFT break down?**


Exploring the limits of chiral EFT



CD, Melendez *et al.*, PRC **102**, 054315

Bayesian inference of the in-medium breakdown scale


But: at what density does chiral EFT break down?



CD, Han, Lattimer *et al.*, PRC **103**, 045808
CD, Han, and Reddy, PRC **105**, 035808

derived **bounds on the neutron star radius** (and sound speed) assuming chiral EFT is valid up to a given critical density (here: $2n_0$) could already be challenged by NICER

$R_{2.0} = (11.4 - 16.1)$ km Riley *et al.*, AJL **918**, L27
Miller *et al.*, AJL **918**, L28




Han & Prakash, APJ **899**, 2
Alford *et al.*, JPG: NPP **46**, 114001

extend EFT EOS at n_m to linear EoS with finite discontinuity (softening)

continuous match sets upper bound

use **lower limit on M_{\max}** from observation to adjust $\Delta\epsilon$ and constrain R_{\min}


Exploring the limits of chiral EFT



CD, Melendez *et al.*, PRC **102**, 054315

Bayesian inference of the in-medium breakdown scale


But: at what density does chiral EFT break down?



CD, Han, Lattimer *et al.*, PRC **103**, 045808
CD, Han, and Reddy, PRC **105**, 035808

derived **bounds on the neutron star radius** (and sound speed) assuming chiral EFT is valid up to a given critical density (here: $2n_0$) could already be challenged by NICER

$R_{2.0} = (11.4 - 16.1)$ km Riley *et al.*, AJL **918**, L27
Miller *et al.*, AJL **918**, L28



Han & Prakash, APJ **899**, 2
Alford *et al.*, JPG: NPP **46**, 114001


extend EFT EOS at n_m to linear EoS with finite discontinuity (softening)

continuous match sets upper bound

use **lower limit on M_{max}** from observation to adjust $\Delta\epsilon$ and constrain R_{min}

Direct astrophysical tests at supranuclear densities

OHIO
UNIVERSITY



CD, Furnstahl *et al.*, PRL 125, 202702

see also: Essick *et al.*, PRC 102, 055803




Neutron star observations could be used for:
 Model checking & selection of chiral interactions
 Constraints on coupling constants in nuclear forces


$$P(n = 0.32 \text{ fm}^{-3}) \approx \begin{cases} 20 \pm 6 \text{ MeV fm}^{-3} & \text{MBPT: nonlocal} \\ 15 \pm 5 \text{ MeV fm}^{-3} & \text{QMC: local } V_{E,1} \end{cases}$$

Direct astrophysical tests at supranuclear densities


OHIO
UNIVERSITY



CD, Furnstahl *et al.*, PRL **125**, 202702



see also: Essick *et al.*, PRC **102**, 055803




CD, Melendez *et al.*, PRC **102**, 054315

Neutron star observations could be used for:
Model checking & selection of chiral interactions
Constraints on coupling constants in nuclear forces

EFT truncation error is *highly* correlated

$$P(n = 0.32 \text{ fm}^{-3}) \approx \begin{cases} 20 \pm 6 \text{ MeV fm}^{-3} & \text{MBPT: nonlocal} \\ 15 \pm 5 \text{ MeV fm}^{-3} & \text{QMC: local } V_{E,1} \end{cases}$$

New predictions in SNM at intermediate densities




Leonhardt, Pospiech, Schallmo, Braun, CD,
Hebeler, and Schwenk, PRL 125, 142502

Functional Renormalization Group (based on QCD action):
ab initio constraints at intermediate densities ($\sim 3 - 10n_0$)


suggests that the different density regions can be
straightforwardly combined

for neutron star matter, see: Braun & Schallmo; arXiv:2204.00358



BAND Manifesto,
Phillips, Furnstahl, Heinz et al.,
JGP: NP 48 07200


New predictions in SNM at intermediate densities



Leonhardt, Pospiech, Schallmo, Braun, CD,
Hebeler, and Schwenk, PRL **125**, 142502


Functional Renormalization Group (based on QCD action):
ab initio constraints at intermediate densities ($\sim 3 - 10n_0$)
suggests that the different density regions can be
straightforwardly combined

for neutron star matter, see: Braun & Schallmo; arXiv:2204.00358



Huth, Wellenhofer, and Schwenk, PRC **103**, 025803

remarkable consistency between theory
predictions, experiment, and astrophysics



Chiral Effective Field Theory and the High-Density Nuclear Equation of State

Annual Review of Nuclear and Particle Science

Vol. 71:403-432 (Volume publication date September 2021)

First published as a Review in Advance on July 6, 2021

<https://doi.org/10.1146/annurev-nucl-102419-041903>



C. Drischler,^{1,2,3} J.W. Holt,⁴ and C. Wellenhofer^{5,6}

¹Department of Physics, University of California, Berkeley, California 94720, USA

²Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

³Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA; email: drischler@frib.msu.edu

⁴Cyclotron Institute and Department of Physics and Astronomy, Texas A&M University, College Station, Texas 77843, USA

⁵Institut für Kernphysik, Technische Universität Darmstadt, 64289 Darmstadt, Germany

⁶ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, 64291 Darmstadt, Germany

<> Full Text HTML

Download PDF

Article Metrics

Reprints | Download Citation | Citation Alerts

Keywords:

Chiral EFT | neutron stars | MBPT
nuclear matter at zero and finite temperature
Bayesian uncertainty quantification
recent neutron star observations


see also in the same journal:
James Lattimer, Annu. Rev. Nucl. Part. Sci. 71, 433

Open Access

Take-away points

multi-messenger
nuclear precision
FRIB

} era




unique opportunity to obtain a
fundamental understanding of
strongly interacting matter, with
great **potential for discoveries**

- 1 Upcoming observational (and experimental) campaigns will provide **stringent constraints** on the properties of neutron stars.
- 2 Chiral EFT enables **microscopic predictions** of nuclear matter (and nuclei) with **quantified uncertainties** to interpret these empirical constraints.
- 3 Automated MBPT: efficient EOS calculations across a wide range of densities, isospin asymmetries, and temperatures, as well as nuclear interactions.
- 4 Bayesian methods: powerful tools for quantifying & propagating **correlated uncertainties** in EFT-based calculations (*model checking* is important).

Many thanks to:

R. Furnstahl S. Han J. W. Holt J. Lattimer K. McElvain
J. Melendez D. Phillips M. Prakash S. Reddy C. Wellenhofer T. Zhao



BUQEYE Collaboration